

Fractional Integration of Multivariable H-function and M-series via Pathway Model

Deepti Arora

Assistant Professor, Department of Mathematics, Jaipur Engineering College, Jaipur

Abstract

In this paper, We speak the partial integration operator configuration formula referred to as Nair Because Channel with the very last fabricated from the multivariate characteristic H and Generalized M series. Additionally, some exciting outcomes are received in phrases of natural consequences, which include the Meijer G function, the Mittag-Leffler feature, Bessel function and terrific longitudinal characteristic. We also obtained two essential formulations of formulas concerning left-fractional elements Riemann-Liouville integrated in segment 5. All effects acquired right here are of a widespread nature and may supply a positive number of results in special function theory.

Keywords: Pathway fractional integral operator, Multivariable H-function, Generalized M-series, Mittag-Leffler function, Bessel-function, Hypergeometric function.

Introduction

Integrated operators with totally different special functions have found vital importance and applications within the numerous subfields of mathematical analysis applied. In recent years, half calculus has become one in every of the quickest growing fields of science and arithmetic. Over the course of 4 decades, variety of researchers like Choi et al. [4], Saijo [18,19], Khan et al.. [3], Kryakova [11,12], Kilbas [23,24], Machado et al. [13,14], Calla [5,6], Calla and Saxena [8] examined comprehensive the various properties, applications, and accessories of the many totally different pressure gauges with the goal of partial integration. several half calculus applications may be found, for instance, in disturbances and fluid dynamics, shape mechanics, random random dynamics, physics and controlled nuclear fusion, nonlinear biological systems, uranology.

Let $f(x) = L(a, b), \alpha \in \mathbb{C}, \Re(\alpha) > 0$, then left sided Riemann-Liouville fractional integral operator defined as

$$(I_{0+}^{\alpha} f)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt \quad (1)$$

For more details, we refer to Kiryakova [11], Kilbas-Srivastava-Trujillo [20] and Samko-Kilbas-Marichev [21] etc.

If $f(t)$ is replaced by $t^{\gamma} f(t)$ in (1), the above operator turns out to be Erdélyi-Kober fractional integral; if it is replaced by ${}_2F_1\left(\eta + \beta, -\gamma; \eta; 1 - \frac{t}{x}\right) f(t)$, then (1) takes the form of Saigo hypergeometric fractional integral

$$\frac{\Gamma(\eta)}{x^{-\eta-\beta}} I_{0+}^{\eta, \beta, \gamma} f(x) = \int_0^x (x-t)^{\eta-1} {}_2F_1\left(\eta + \beta, -\gamma; \eta; 1 - \frac{t}{x}\right) f(t) dt$$

Many other operators generalized fractional calculus can be obtained if on the place of $f(t)$ one takes $\phi(t)$ as it is done in Kiryakova [11] for a Fox's H-function $\phi(t) = H_{m,m}^{m,0}(t)$.

The pathway fractional integration operator, as an extension of (1), is defined as follows [2, p.239]:

$$\left(P_{0+}^{(\eta, \lambda, d)} f\right)(x) = x^{\eta} \int_0^{\left[\frac{x}{d(1-\lambda)}\right]} \left[1 - \frac{d(1-\lambda)t}{x}\right]^{\frac{\eta}{1-\lambda}} f(t) dt, \quad (2)$$

where $f(x) \in L(a, b)$, $\eta \in \mathbb{C}$, $\Re(\eta) > 0$, $d > 0$ and 'pathway parameters' $\lambda < 1$.

The pathway model is introduced by Mathai [25] and studied further by Mathai and Haubold [26, 27]. For real scalar λ , the pathway model for scalar random variables is represented by the following probability density function (p.d.f.):

$$f(x) = c|x|^{\gamma-1} [1 - d(1-\lambda)|x|^{\delta}]^{\frac{\beta}{1-\lambda}}, \quad (3)$$

$-\infty < x < \infty$, $\delta > 0$, $\beta \geq 0$, $1 - d(1-\lambda)|x|^{\delta} > 0$, $\gamma > 0$, where c is the normalization constant is as follows:

$$c = \frac{1}{2} \frac{\delta [d(1-\lambda)]^{\frac{\gamma}{\delta}} \Gamma\left(\frac{\gamma}{\delta} + \frac{\beta}{1-\lambda} + 1\right)}{\Gamma\left(\frac{\gamma}{\delta}\right) \Gamma\left(\frac{\beta}{1-\lambda} + 1\right)}, \text{ for } \lambda < 1, \quad (4)$$

$$= \frac{1}{2} \frac{\delta [d(1-\lambda)]^{\frac{\gamma}{\delta}} \Gamma\left(\frac{\beta}{1-\lambda}\right)}{\Gamma\left(\frac{\gamma}{\delta}\right) \Gamma\left(\frac{\beta}{1-\lambda} - \frac{\gamma}{\delta}\right)}, \text{ for } \frac{1}{\lambda-1} - \frac{\gamma}{\delta} > 0, \lambda > 1, \quad (5)$$

$$= \frac{1}{2} \frac{\delta (\lambda\beta)^{\frac{\gamma}{\delta}}}{\Gamma\left(\frac{\gamma}{\delta}\right)} \text{ for } \lambda \rightarrow 1. \quad (6)$$

For $\lambda < 1$, it is a finite range density with $1 - d(1-\lambda)|x|^{\delta} > 0$ and (3) remains in the extended generalized type-1 beta family. The pathway density in (3) for $\lambda < 1$, includes the extended type-1 beta density, the triangular density, the uniform density and many other p.d.f.

For $\lambda > 1$, writing $1 - \lambda = -(\lambda - 1)$, we have

$$f(x) = c|x|^{\gamma-1} [1 + d(\lambda - 1)|x|^{\delta}]^{-\frac{\beta}{\lambda-1}}, \quad (7)$$

$-\infty < x < \infty$, $\delta > 0$, $\beta \geq 0$, $\lambda > 1$, which is extended generalized type-2 beta models for real x . It includes the type-2 beta density, the F density, the Cauchy density and many more.

Here we consider only the case of pathway parameters $\lambda < 1$. For $\lambda \rightarrow 1$ both (3) and (7) take the exponential form, since

$$\begin{aligned} \lim_{\lambda \rightarrow 1} c|x|^{\gamma-1} [1 - d(1-\lambda)|x|^{\delta}]^{\frac{\eta}{1-\lambda}} &= \lim_{\lambda \rightarrow 1} c|x|^{\gamma-1} \left[1 - d \eta |x|^{\delta} \frac{(1-\lambda)}{\eta}\right]^{\frac{\eta}{1-\lambda}} \\ &= c|x|^{\gamma-1} \exp(-d \eta |x|^{\delta}). \end{aligned} \quad (8)$$

This includes the generalized Gamma-, the Weibull-, the Chi-square, the Laplace-, Maxwell-Boltzmann and other related densities. Therefore, the operator introduced in this paper can be related and applicable to a wide variety of statistical density.

When $\lambda \rightarrow 1_-$, $\left[1 - \frac{d(1-\lambda)t}{x}\right]^{\frac{\eta}{1-\lambda}} \rightarrow e^{-d\eta t}$. Then, the operators (1) reduces to the Laplace integral transform of f with the parameters $\frac{d\eta}{x}$:

$$(P_{0+}^{(\eta,1)} f)(x) = x^\eta \int_0^\infty e^{-\frac{d\eta}{x}t} f(t) dt = x^\eta L_f\left(\frac{d\eta}{x}\right). \quad (9)$$

Mathematical Prerequisites

The multivariable H-function is defined in terms of multiple Mellin-Barnes type contour integral as [1].

$$H[z_1, \dots, z_r] = H_{p,q; p_1, q_1, \dots, p_r, q_r}^{0, n; m_1, n_1, \dots, m_r, n_r} \left[\begin{matrix} z_1 \\ \vdots \\ z_r \end{matrix} \left| \begin{matrix} (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,p} : (c_j^{(1)}; \gamma_j^{(1)})_{1,p_1} \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)})_{1,q} : (d_j^{(1)}; \delta_j^{(1)})_{1,q_1} \\ \vdots \\ (c_j^{(r)}; \gamma_j^{(r)})_{1,p_r} \\ (d_j^{(r)}; \delta_j^{(r)})_{1,q_r} \end{matrix} \right. \right] \quad (10)$$

$$H[z_1, \dots, z_r] = \frac{1}{(2\pi\omega)^r} \int_{\mathcal{L}_1} \dots \int_{\mathcal{L}_r} \Theta(\xi_1 \dots \xi_r) \left\{ \prod_{i=1}^r \varphi_i(\xi_i) z_i^{\xi_i} \right\} d\xi_1 \dots d\xi_r$$

where

$$\Theta(\xi_1 \dots \xi_r) = \frac{\left[\prod_{j=1}^n \Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(i)} \xi_i) \right]}{\left[\prod_{j=n+1}^p \Gamma(a_j - \sum_{i=1}^r \alpha_j^{(i)} \xi_i) \right] \left[\prod_{j=1}^q \Gamma(1 - b_j + \sum_{i=1}^r \beta_j^{(i)} \xi_i) \right]} \quad (11)$$

$$\varphi_i(\xi_i) = \frac{\left[\prod_{j=1}^{m_i} \Gamma(d_j^{(i)} - \delta_j^{(i)} \xi_i) \right] \left[\prod_{j=1}^{n_i} \Gamma(1 - c_j^{(i)} + \gamma_j^{(i)} \xi_i) \right]}{\left[\prod_{j=n_i+1}^{p_i} \Gamma(c_j^{(i)} - \gamma_j^{(i)} \xi_i) \right] \left[\prod_{j=m_i+1}^{q_i} \Gamma(1 - d_j^{(i)} + \delta_j^{(i)} \xi_i) \right]} \quad (12)$$

For $i = 1, \dots, r$ and $\omega = \sqrt{-1}$, $\mathcal{L}_i$ represents the contour which start at the point $\tau_i - \omega\infty$ and goes out the point $\tau_i + \omega\infty$ with $\tau_i \in \mathbb{R} = (-\infty, \infty)$, $i = 1, \dots, r$ such that all the poles of $\Gamma(d_j^{(i)} - \delta_j^{(i)} \xi_i)$, $j = 1, \dots, m_i$; $i = 1, \dots, r$

are separated from those of $\Gamma(1 - c_j^{(i)} + \gamma_j^{(i)} \xi_i)$, $j = 1, \dots, n_i$; $i = 1, \dots, r$ and

$\Gamma(1 - a_j + \sum_{i=1}^r \alpha_j^{(i)} \xi_i)$, $j = 1, \dots, n$. Here, the integers n, p, q, m_i, n_i, p_i and q_i , satisfy the inequalities

$0 \leq n \leq p$; $q \geq 0$, $0 \leq m_i \leq q_i$ and $0 \leq n_i \leq p_i$,

$i = 1, \dots, r$. Further we suppose that the parameters

$$\begin{aligned} a_j, j = 1, \dots, p; c_j^{(i)}, j = 1, \dots, p_i; i = 1, \dots, r \\ b_j, j = 1, \dots, q; d_j^{(i)}, j = 1, \dots, q_i; i = 1, \dots, r \end{aligned} \quad (13)$$

are complex numbers and the related coefficients

$$\begin{aligned} \alpha_j^{(i)}, j = 1, \dots, p; i = 1, \dots, r; \gamma_j^{(i)}, j = 1, \dots, p_i; i = 1, \dots, r \\ \beta_j^{(i)}, j = 1, \dots, q; i = 1, \dots, r; \delta_j^{(i)}, j = 1, \dots, q_i; i = 1, \dots, r \end{aligned} \quad (14)$$

are positive real numbers.

We observe that for $n = m = p = q = 0$, the multivariable H-function breaks up in to product of r H -function and consequently there holds the following result [7].

$$\begin{aligned} H[z_1, \dots, z_r] &= H_{0,0;p_1,q_1;\dots;p_r,q_r}^{0,0;m_1,n_1;\dots;m_r,n_r} \left[\begin{matrix} z_1 \\ \vdots \\ z_r \end{matrix} \middle| \begin{matrix} -; (c_j^{(1)}; \gamma_j^{(1)})_{1,p_1}; \dots; (c_j^{(r)}; \gamma_j^{(r)})_{1,p_r} \\ -; (d_j^{(1)}; \delta_j^{(1)})_{1,q_1}; \dots; (d_j^{(r)}; \delta_j^{(r)})_{1,q_r} \end{matrix} \right] \\ &= \prod_{i=1}^r H_{p_i,q_i}^{m_i,n_i} \left[z_i \middle| \begin{matrix} (c_j^{(i)}; \gamma_j^{(i)})_{1,p_i} \\ (d_j^{(i)}; \delta_j^{(i)})_{1,q_i} \end{matrix} \right] \end{aligned} \quad (15)$$

If we consider $n = m = p = q = 0$ and $r = 1$ then the multivariable H-function converts in to as single variable H-function and it becomes

$$H[z] = H_{p,q}^{m,n}[z] = H_{p,q}^{m,n} \left[z \middle| \begin{matrix} (a_p; \alpha_p) \\ (a_q; \beta_q) \end{matrix} \right] = \frac{1}{2\pi\omega} \int_{\mathcal{L}} \Theta(\xi) z^{-\xi} d\xi, \quad (16)$$

where

$$\Theta(\xi) = \frac{\prod_{j=1}^m \Gamma(b_j + \beta_j \xi) \prod_{j=1}^n \Gamma(1 - a_j - \alpha_j \xi)}{\prod_{j=m+1}^q \Gamma(1 - b_j - \beta_j \xi) \prod_{j=n+1}^p \Gamma(a_j + \alpha_j \xi)} \quad (17)$$

$m, n, p, q \in N_0$ with $1 \leq n \leq p, 1 \leq m \leq q$, $\alpha_j, \beta_j \in \mathbb{R}_+$ and $a_j, b_j \in \mathbb{R}$ ($i = 1, \dots, p; j = 1, \dots, q$) and $\mathcal{L}$ is a suitable contour which separating the poles of $\Gamma(b_j + \beta_j \xi)$ from poles $\prod_{j=1}^n \Gamma(1 - a_j - \alpha_j \xi)$.

Wright [22] defined generalized hypergeometric function by means of the series representation in the form

$${}_p\psi_q(z) = {}_p\psi_q \left[\begin{matrix} (a_1, A_1), \dots, (a_p, A_p) \\ (b_1, B_1), \dots, (b_q, B_q) \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + nA_i) z^n}{\prod_{j=1}^q \Gamma(b_j + nB_j) n!} \quad (18)$$

where $z, a_j, b_j \in \mathbb{C}, A_i, B_j \in \mathbb{R}_+$ ($i = 1, \dots, p; j = 1, \dots, q$),

$$\sum_{j=1}^q B_j - \sum_{i=1}^p A_i > -1 \quad (19)$$

Sharma and Jain [9] introduced the generalized M-series as the function defined by means of the power series:

$$\begin{aligned} {}^{\alpha}_p M_q^{\beta}(z) &= {}^{\alpha}_p M_q^{\beta}(a_1, \dots, a_p; b_1, \dots, b_q; z) = {}^{\alpha}_p M_q^{\beta}\left((a_j)_1^p; (b_j)_1^q; z\right) \\ &= \sum_{n=0}^{\infty} \frac{(a_1)_n, \dots, (a_p)_n}{(b_1)_n, \dots, (b_q)_n} \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad (z, \alpha, \beta \in \mathbb{C}, \Re(\alpha) > 0) \end{aligned} \quad (20)$$

where $(a_j)_n, (b_j)_n$ are known Pochhammer symbols. The series (20) is defined when none of the parameters b_j 's, $j = 1, 2, \dots, q$ is a negative integer or zero; if any numerator parameter a_j is a negative integer or zero, then the series terminates to a polynomial in z . The series in (20) is convergent for all z if $p \leq q$, it is convergent for $|z| < \delta = \alpha^{\alpha}$ if $p = q + 1$ and divergent, if $p > q + 1$. When $p = q + 1$ and $|z| = \delta$, the series can converge on conditions depending on the parameters. Properties of M-series are further studied by Saxena [10], Chouhan and Sarswat [17] etc.

The generalized M-series (20) can be represented as a special case of Fox H-function (16) and Wright generalized hypergeometric function (18), as

$$\begin{aligned} {}^{\alpha}_p M_q^{\beta}\left((a_j)_1^p; (b_j)_1^q; z\right) &= \Lambda_{p+1} \psi_{q+1} \left[\begin{matrix} (a_1, 1), \dots, (a_p, 1), (1, 1); \\ (b_1, 1), \dots, (b_q, 1), (\beta, \alpha); \end{matrix} z \right] \\ &= \Lambda H_{p+1, q+1}^{1, p+1} \left[-z \left| \begin{matrix} (1 - \alpha_j, 1; 1)_1^p, (0, 1) \\ (0, 1), (1 - \beta_j, 1)_1^q, (1 - \beta, \alpha) \end{matrix} \right. \right] \end{aligned} \quad (21)$$

$$\text{where } \Lambda = \frac{\prod_{j=1}^q \Gamma(b_j)}{\prod_{j=1}^p \Gamma(a_j)}.$$

The generalized Mittag-Leffler function, introduced by Prabhakar [32] may be obtain from (21) for $p = q = 1; \alpha = \gamma \in \mathbb{C}; b = 1$, as

$$E_{\alpha, \beta}^{\gamma}(z) = \sum_{m=0}^{\infty} \frac{(\gamma)_m}{\Gamma(\alpha m + \beta)} \frac{z^m}{m!} = \sum_{m=0}^{\infty} \frac{(\gamma)_m}{(1)_m} \frac{z^m}{\Gamma(\alpha m + \beta)} = {}^{\alpha}_1 M_1^{\beta}(1; 1; z) \quad (22)$$

The new generalization Mittag-Leffler function in terms of H-function is defined as

$$E_{\alpha, \beta}^{\gamma}(z) = \frac{1}{\Gamma(\gamma)} H_{1, 2}^{1, 1} \left[-z \left| \begin{matrix} (1 - \gamma, 1) \\ (0, 1), (1 - \beta, \alpha) \end{matrix} \right. \right] \quad (\alpha, \beta, \gamma \in \mathbb{C}; \Re(\alpha) > 0) \quad (23)$$

The present paper is organized as follows. The composition of pathway integral operators (2) with multivariables H -function, Mittag-Leffler, Bessel- function are given in section 3. Section 4 investigates the pathway fractional integration of generalized M-series and finally the further special cases in section 5 and concluding remark are drawn in section 6.

Fractional Integration of H-function via Pathway Model

Recently pathway fractional integral operators involving the various special functions have been considered by many author (see, eq., [29, 30, 31]). Our main result in this section is based on the following assertion giving a composition formula of the pathway fractional integration operator (2) with a power function (see Nair [2, Lemma 1]).

Lemma 1. Let $\eta \in \mathbb{C}, \Re(\eta) > 0, \beta \in \mathbb{C}$ and $\lambda < 1$. If $\Re(\beta) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then we have

$$\left\{P_{0+}^{(\eta, \lambda, d)} [t^{\beta-1}]\right\}(x) = \frac{x^{\eta+\beta} \Gamma(\beta) \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[d(1-\lambda)]^{\beta} \Gamma\left(1 + \frac{\eta}{1-\lambda} + \beta\right)} \quad (24)$$

Now we are in a position to state and prove our main result, which is a composition formula of the pathway fractional integration operator (2) with a finite product of multivariable H-function (10) asserted by the following Theorem.

Theorem 1. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z_i, \eta, \rho \in \mathbb{C} (i = 1, \dots, r), \Re(\eta) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then there holds the following formula:

$$\begin{aligned} & P_{0+}^{(\eta, \lambda, l)} \{t^{\rho-1} H[d^{\mu} t^{\mu} z_1, \dots, d^{\mu} t^{\mu} z_r]\} \\ &= \frac{t^{\eta+\rho} \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[l(1-\lambda)]^{\rho}} H_{p+1, q+1; p_1, q_1; \dots; p_r, q_r}^{0, n+1; m_1, n_1; \dots; m_r, n_r} \left[\begin{matrix} d^{\mu} t^{\mu} z_1 \\ \{l(1-\lambda)\} \\ \vdots \\ d^{\mu} t^{\mu} z_r \\ \{l(1-\lambda)\} \end{matrix} \middle| \begin{matrix} (1-\rho; \mu, \dots, \mu), (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,p} \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)})_{1,q}, \left(-\frac{\eta}{1-\lambda} - \rho; \mu, \dots, \mu\right) \end{matrix} \right. \\ & \left. : (c_j^{(1)}; \gamma_j^{(1)})_{1,p_1}; \dots; (c_j^{(r)}; \gamma_j^{(r)})_{1,p_r} \right. \\ & \left. : (d_j^{(1)}; \delta_j^{(1)})_{1,q_1}; \dots; (d_j^{(r)}; \delta_j^{(r)})_{1,q_r} \right] \quad (25) \end{aligned}$$

Proof. Let $\mathfrak{I}$ be the left-hand side of (25). By applying (10) and using (2) to left hand side of (25), then we have

$$\mathfrak{I} = \frac{1}{(2\pi\omega)^r} \int_{\mathcal{L}_1} \dots \int_{\mathcal{L}_r} \Theta(\xi_1 \dots \xi_r) \left\{ \prod_{i=1}^r \varphi_i(\xi_i) (d^{\mu} z_i)^{\xi_i} \right\} d\xi_1 \dots d\xi_r \times P_{0+}^{(\eta, \lambda, l)} \{t^{\rho+\mu \sum_{i=1}^r \xi_i-1}\}$$

Now we apply Lemma 1 to use (24) with β replaced by $(\rho + \mu \sum_{i=1}^r \xi_i)$ to pathway integral, after a little simplification, we obtain the following result

$$\begin{aligned} & \mathfrak{I} = \frac{1}{(2\pi\omega)^r} \int_{\mathcal{L}_1} \dots \int_{\mathcal{L}_r} \Theta(\xi_1 \dots \xi_r) \left\{ \prod_{i=1}^r \varphi_i(\xi_i) (d^{\mu} z_i)^{\xi_i} \right\} d\xi_1 \dots d\xi_r \\ & \times t^{\eta+\rho+\mu \sum_{i=1}^r \xi_i} \left(\frac{\Gamma(\rho + \mu \sum_{i=1}^r \xi_i) \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[l(1-\lambda)]^{\rho+\mu \sum_{i=1}^r \xi_i} \Gamma\left(\rho + \frac{\eta}{1-\lambda} + \mu \sum_{i=1}^r \xi_i + 1\right)} \right) \end{aligned}$$

then, in view of (10), interpreting the involved Mellin-Barnes contour integrals in term of Multivariable H-function, we readily obtain the right hand side of (25).

Setting $n = p = q = 0$ in (25) is seen to break the multivariable H-function in (15) into an r-times product of H-functions yield an interesting result asserted by the following Corollary 1.

Corollary 1. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z_i, \eta, \rho \in \mathbb{C} (i = 1, \dots, r), \Re(\eta) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then there holds the following formula:

$$P_{0+}^{(\eta, \lambda, l)} \left\{ t^{\rho-1} \prod_{i=1}^r H_{p_i, q_i}^{m_i, n_i} \left[d^\mu t^\mu z_i \left| \begin{matrix} (c_j^{(i)}; \gamma_j^{(i)})_{1, p_i} \\ (d_j^{(i)}; \delta_j^{(i)})_{1, q_i} \end{matrix} \right. \right] \right\}$$

$$= \frac{t^{\eta+\rho} \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[l(1-\lambda)]^\rho} H_{1,1;\{p_i, n_i\}}^{0,1;\{m_i, q_i\}} \left[\frac{d^\mu t^\mu z_i}{\{l(1-\lambda)\}} \left| \begin{matrix} (1-\rho; \mu, \dots, \mu): (c_j^{(i)}; \gamma_j^{(i)})_{1, p_i} \\ \left(-\frac{\eta}{1-\lambda} - \rho; \mu, \dots, \mu\right): (d_j^{(i)}; \delta_j^{(i)})_{1, q_i} \end{matrix} \right. \right]$$

(26)

If we observe that for $n = p = q = 0$ and $r = 1$, the multivariable H-function reduces in to single variable H-function and consequently there holds the following Corollary.

Corollary 2. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z, \eta, \rho \in \mathbb{C}, \Re(\eta) > 0, \Re(\rho) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then we obtain

$$P_{0+}^{(\eta, \lambda, l)} \{ t^{\rho-1} H_{p,q}^{m,n} (d^\mu t^\mu z) \}$$

$$= \frac{t^{\eta+\rho} \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[l(1-\lambda)]^\rho} \times H_{p+1,q+1}^{m,n+1} \left[\frac{d^\mu t^\mu z}{\{l(1-\lambda)\}} \left| \begin{matrix} (1-\rho, \mu), (\alpha_p, \alpha_p) \\ \left(-\frac{\eta}{1-\lambda} - \rho, \mu\right), (b_q, \beta_q) \end{matrix} \right. \right] \quad (27)$$

Setting $\alpha_i = \beta_j = 1$ in the Corollary 2, yields the following result.

Corollary 3. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z, \eta, \rho \in \mathbb{C}, \Re(\eta) > 0, \Re(\rho) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then we obtain

$$P_{0+}^{(\eta, \lambda, l)} \{ t^{\rho-1} G_{p,q}^{m,n} (d^\mu t^\mu z) \}$$

$$= \frac{t^{\eta+\rho} \Gamma\left(1 + \frac{\eta}{1-\lambda}\right)}{[l(1-\lambda)]^\rho} \times G_{p+1,q+1}^{m,n+1} \left[\frac{d^\mu t^\mu z}{\{l(1-\lambda)\}} \left| \begin{matrix} (1-\rho, \mu), (a_i)_1^p \\ \left(-\frac{\eta}{1-\lambda} - \rho, \mu\right), (b_j)_1^q \end{matrix} \right. \right] \quad (28)$$

Interestingly,

on

setting $m = 1, p = 1, n = 1, q = 2, \mu = 1, z = -Y, \alpha_1 = 1 - \gamma, \alpha_1 = 1, b_1 = 0, \beta_1 = 1, b_2 = 1 - \beta, \beta_2 = \alpha$, then Corollary 2 and consequently there holds the following result.

Corollary 4. Let $\lambda < 1, d > 0, l > 0$, the parameters $\alpha, \beta, \gamma, \eta, \rho \in \mathbb{C}, \Re(\alpha) > 0, \Re(\eta) > 0, \Re(\rho) > 0$ and $\Re\left(\frac{\eta}{1-\lambda}\right) > -1$, then we have

$$\begin{aligned}
 & P_{0+}^{(\eta, \lambda, l)} \left\{ t^{\rho-1} E_{\alpha, \beta}^{\gamma} (Ytd) \right\} \\
 &= \frac{t^{\eta+\rho} \Gamma \left(1 + \frac{\eta}{1-\lambda} \right)}{[l(1-\lambda)]^{\rho} \Gamma(\gamma)} \times H_{2,3}^{1,2} \left[\begin{matrix} (1-\rho, 1), (1-\gamma, 1) \\ (0, 1), \left(-\frac{\eta}{1-\lambda} - \rho, 1 \right), (1-\beta, \alpha) \end{matrix} ; \frac{-Ytd}{[l(1-\lambda)]} \right] \quad (29)
 \end{aligned}$$

Further, if we set $m = 1, n = 0, p = 0, q = 0, \mu = 2, b_1 = 0, \beta_1 = 1, b_2 = v, \beta_2 = 1, \rho = \sigma + \vartheta$ in Corollary 2, then we obtain the following Corollary.

Corollary 5. Let the condition of Corollary 2 be satisfied then the equation (26) reduces to the following result.

$$\begin{aligned}
 & P_{0+}^{(\eta, \lambda, l)} \{ t^{\rho-1} J_v(dtz) \} \\
 &= \left(\frac{zd}{2} \right)^v \frac{t^{\eta+\sigma+\vartheta} \Gamma \left(1 + \frac{\eta}{1-\lambda} \right)}{[l(1-\lambda)]^{\sigma+\vartheta}} \times {}_1\psi_2 \left[\begin{matrix} (\sigma + \vartheta, 2) \\ (\vartheta + 1, 1), \left(1 + \frac{\eta}{1-\lambda} + \sigma + \vartheta, 2 \right) \end{matrix} ; \frac{-d^2 t^2 z^2}{4[l(1-\lambda)]} \right] \quad (30)
 \end{aligned}$$

where $J_v(z)$ is the ordinary Bessel function of first kind (Olver [15]).

Corollary 6. Let the condition of corollary 2 be satisfied and on setting $m = 1, n = 1, p = 1, q = 2, \alpha_1 = 1 - \alpha, \alpha_1 = 1, b_1 = 0, \beta_1 = 0, b_2 = 1 - c, \mu = 1, \beta_2 = 1, z = -Y$, then equation (26) reduces to the following formula:

$$\begin{aligned}
 & P_{0+}^{(\eta, \lambda, l)} \{ t^{\rho-1} {}_1F_1(a; c; dtY) \} \\
 &= \frac{t^{\eta+\rho} \Gamma \left(1 + \frac{\eta}{1-\lambda} \right) \Gamma(\rho)}{[l(1-\lambda)]^{\rho} \Gamma \left(\frac{\eta}{1-\lambda} + \rho + 1 \right)} \times {}_2F_2 \left[\begin{matrix} a, (\rho) \\ c, \left(\frac{\eta}{1-\lambda} + \rho + 1 \right) \end{matrix} ; \frac{Ytd}{[l(1-\lambda)]} \right] \quad (31)
 \end{aligned}$$

Provided $\Re(\eta) > 0, \Re(\rho) > 0$ and $\Re \left(\frac{\eta}{1-\lambda} \right) > -1$.

Fractional Integration of Generalized M-series via Pathway Model

In this section we consider composition of the pathway fractional integral $P_{0+}^{(\eta, \lambda, l)}$ given by (2) with the generalized M-series (20).

Theorem 2. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z, \alpha, \beta, \gamma, \eta, \rho \in \mathbb{C}$ and $\Re(\eta) > 0$,

$\Re(\alpha) > 0, \Re(\rho) > 0, \Re \left(\frac{\eta}{1-\lambda} \right) > -1$, then the following formula holds:

$$\begin{aligned}
 & P_{0+}^{(\eta, \lambda, l)} \left\{ t^{\rho-1} {}_{\alpha}M_q^{\beta} (d^{\mu} t^{\mu} z) \right\} = \frac{t^{\eta+\rho} \Gamma \left(1 + \frac{\eta}{1-\lambda} \right) \sum_{j=1}^q \Gamma(b_j)}{[l(1-\lambda)]^{\rho} \sum_{i=1}^p \Gamma(a_i)} \\
 & \times H_{p+2, q+3}^{1, p+2} \left[\begin{matrix} (1-\rho, \mu), (1-a_i)_{i=1}^p, (0, 1) \\ -\frac{d^{\mu} t^{\mu} z}{[l(1-\lambda)]}, \left(-\frac{\eta}{1-\lambda} - \rho, \mu \right), (0, 1), (1-\beta, \alpha), (1-b_j)_{j=1}^q \end{matrix} \right] \quad (32)
 \end{aligned}$$

The converges condition for the validity (32) are as follows

(i) $0 \leq m \leq q, 0 \leq n \leq p$ for $a_i, b_j \in \mathbb{C}$ and for $\alpha_i, \beta_j \in \mathbb{R}_+ = (0, \infty)$ ($i = 1, \dots, p; j = 1, \dots, q$).

(ii) $\mathcal{L} = \mathcal{L}_{i\gamma\infty}$ is a contour starting at the point $\gamma - i\infty$ and going to $\gamma + i\infty$ where $\gamma \in \mathbb{R} (-\infty, \infty)$, such that all the poles of $\Gamma(b_j + \beta_j \xi), j = 1, \dots, m$ are separated from those $\Gamma(1 - a_j - \alpha_j \xi), i = 1, \dots, n$.

The integral converges if $\alpha > 0, |\arg z| < \frac{1}{2}\pi\alpha, \alpha \neq 0$ and $\sigma = \sum_{j=1}^n \alpha_j - \sum_{j=n+1}^p \alpha_j + \sum_{j=1}^m \beta_j - \sum_{j=m+1}^q \beta_j$.

Proof. For convenience, let the left hand side of (32) be denoted by $\square$. Applying (20) and using (2) to left hand side of (32), then we have

$$\square = \sum_{m=0}^{\infty} \frac{(a_1)_m \dots (a_p)_m}{(b_1)_m \dots (b_q)_m} \frac{(d^\mu z)^m}{\Gamma(\alpha m + \beta)} \times \left[P_{0+}^{(\eta, \lambda, l)}(t^{\rho + \mu m - 1}) \right]$$

Now we apply Lemma 1 to use (24) with β replaced by $(\rho + \mu m)$ to pathway integral, after a little simplification, we obtain

$$\square = \frac{t^{\eta + \rho} \Gamma\left(1 + \frac{\eta}{1 - \lambda}\right) \sum_{j=1}^q \Gamma(b_j)}{[l(1 - \lambda)]^\rho \sum_{i=1}^p \Gamma(a_i)} \sum_{m=0}^{\infty} \frac{\Gamma(a_i + m) \Gamma(\rho + \mu m)}{\Gamma(b_j + m) \Gamma(\alpha m + \beta)}$$

$$\times \frac{1}{\Gamma\left(\frac{\eta}{1 - \lambda} + \rho + \mu m + 1\right)} \left[\frac{d^\mu t^\mu z}{l(1 - \lambda)} \right]^m$$

which, upon using definition of H-function (16), yields (32).

Several illustrative examples of Theorem 7 involving appropriately chosen special values of parameters can also be derived fairly easily.

Further Special Cases

By setting $\lambda = 0, l = 1$ and $\eta \rightarrow \eta - 1$ in (25) and (32) respectively, and applying the following easily derivable relation:

$$\left(P_{0+}^{(\eta-1, 0, 1)} f \right) (t) = \int_0^t (t - \tau)^{\eta-1} f(\tau) d\tau = \Gamma(\eta) (I_{0+}^\eta f)(t), \quad (\square(\eta) > 0)$$

We obtain two fractional integral formulas involving left-sided Riemann-Liouville fractional integral operator stated in the next Corollaries below.

Corollary 7. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z_i, \eta, \rho \in \mathbb{R} (i = 1, \dots, r), \square(\eta) > 0, \square(\rho) > 0$ and

$\square\left(\frac{\eta}{1 - \lambda}\right) > -1$, then the following relation

$$I_{0+}^\eta \{t^{\rho-1} H[d^\mu t^\mu z_1, \dots, d^\mu t^\mu z_r]\}$$

$$= t^{\rho + \eta - 1} \Gamma(\eta) H_{p+1, q+1; p_1, q_1; \dots; p_r, q_r}^{0, n+1; m_1, n_1; \dots; m_r, n_r} \left[\begin{matrix} (1 - \rho; \mu, \dots, \mu), (a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1, p} \\ (b_j; \beta_j^{(1)}, \dots, \beta_j^{(r)})_{1, q}, (1 - \eta - \rho; \mu, \dots, \mu) \end{matrix} \right]$$

$$\begin{aligned} &: \left(c_j^{(1)}; \gamma_j^{(1)} \right)_{1,p_1}; \dots; \left(c_j^{(r)}; \gamma_j^{(r)} \right)_{1,p_r} \left[\begin{matrix} d^\mu t^\mu z_1 \\ \vdots \\ d^\mu t^\mu z_r \end{matrix} \right] \\ &: \left(d_j^{(1)}; \delta_j^{(1)} \right)_{1,q_1}; \dots; \left(d_j^{(r)}; \delta_j^{(r)} \right)_{1,q_r} \end{aligned} \quad (33)$$

holds.

Corollary 8. Let $\lambda < 1, \mu > 0, d > 0, l > 0$, the parameters $z, \alpha, \beta, \gamma, \eta, \rho \in \mathbb{C}$ and $\Re(\eta) > 0$,

$\Re(\alpha) > 0, \Re(\rho) > 0, \Re\left(\frac{\eta}{1-\lambda}\right) > -1$. Then the following formula relation:

$$\begin{aligned} I_{0+}^\eta \left\{ t^{\rho-1} {}_pM_q^\beta(d^\mu t^\mu z) \right\} &= t^{\eta+\rho-1} \Gamma(\eta) \frac{\sum_{j=1}^q \Gamma(b_j)}{\sum_{i=1}^p \Gamma(a_i)} \\ &\times H_{p+2,q+3}^{1,p+2} \left[-d^\mu t^\mu z \left| \begin{matrix} (1-\rho, \mu), (1-a_i)_1^p, (0,1) \\ (1-\eta-\rho, \mu), (0,1), (1-\beta, \alpha), (1-b_j)_1^q \end{matrix} \right. \right] \end{aligned} \quad (34)$$

holds.

Remark. It is noted that if we set $\lambda = 0, l = 1$ and $f(t)$ is replaced by ${}_2F_1\left(\eta + \beta, -\gamma; \eta; 1 - \frac{t}{x}\right) f(t) dt$, (2) yields the Saigo fractional integral operator. Thus we can obtain the generalizations of left-sided fractional integrals, like Saigo, Erdélyi-Kober (see [28]; see also [20]), and so on, by suitable substitutions. Therefore, the result presented here easily shown to be converted to those corresponding to the above well known fractional operators.

Concluding Remarks

We conclude this analysis by noting that the results obtained here are of a general and helpful nature to derive multiple integral formulas within the theory of integrated micro-channel equation. during this article, we have a tendency to conferred the syntax for aliquot integral for a mechanical phenomenon that has a operate Fox H and M series. The leads to this text are vital and you'll realize an application in resolution differential equations. From an aliquot arrangement that arises in theory from special functions. Associate in nursing integral channel operator with many special functions given as special cases to our main findings.

References

1. H.M. Srivastava and R.Panda, Some bilateral equation generating function for a class of generalized hypergeometric polynomials, J.Reine.Angew. Math., 283/284 (1976), 265-274.
2. S.S. Nair, Pathway fractional integration operator, Fract.Calc. appl. Anal., Vol. 12(3) (2009), 237-252.
3. A.M. Khan, R.K. Kumbhat, A.Chuhan and A.Alaria, Generalized fractional integral operators and M-series, Hindawi Publishing Corporation, Volume 2016, Article ID 2872185, 10 pages.
4. J.Choi, P.Agarwal and S.Jain, Certain fractional integral operators and extended generalization Gauss hypergeometric functions, KYUNGPOOK Math. J., Vol. 55 (2015), 695-703.
5. S.L. Kalla, Integral operators involving Fox's H- function I, Acta Maxicana Cienc. Tecn. , Vol.3 (1969), 117-122.
6. S.L Kalla, Integral operators involving Fox's H- function II, Acta Maxicana Cienc. Tecn. , Vol.7 (1969), 72-79.
7. R.K. Saxena, On the H-function of n-variables, Kyungpook Math. J., vol. 17 (1977), 221-226.

8. S.L. Kalla and R.K. Saxena, Integral operators involving hyper geometric functions, *Math.Z.*, Vol.108 (1969), 231-234.
9. M.Sharma and R.Jain, A note on a generalized M-series as a special function of fractional calculus, *Frac. Calc. Appl. Anl.*, Vol.12 (4), (2009), 449-452.
10. R.K. Saxena, A remark on a paper on M-series, *Fract. Calc. Appl. Anal.*, Vol.12 (1) (2009) , 109-110.
11. V. Kiryakova, Generalized fractional calculus and applications, Longman Scientific & Tech. Essex. (1994).
12. V. Kiryakova, A brief story about the operators of the generalized fractional calculus, *Fract, Calc. Appl.*, Vol.12 (2) (2008), 203-220.
13. J.T. Machado, V. Kiryakova and F.Mainardi, Recent history of fractional calculus communications in Nonlinear science and Numerical Simulation (Elsevier), Vol.16 (2011) 1140-1153; doi :10.1016/j.cnsns.2010.05.027.=
14. J.T. Machado, V. Kiryakova and F.Mainardi, A poster about the recent history of fractional calculus, *Fract . Calc. Appl. Anal.* Vol.13 (3) (2010), 329-334.
15. F.N.L Olver, D.M. Lozier, R.F. Boisvert and C.W. Clark, NIST Hand book of Mathematical Function, Cambridge University Prees, Cambridge 2010.
16. A.C.Mc Bride, Fractional Power of a class of ordinary differential operators, *Proc. London Math. Soc.*, Vol.45 (3) (1982), 519-546.
17. A. Chouhan and S. Saraswat, Certain properties of fractional calculus operators associated with M-series, *Scientia : Series A : Mathematical Sciences* 22 (2012), 25-30 .
18. M. Saigo, A remark on integral operators involving the Gauss hypergeometric functions, *Math. Rep. Kyushu Univ.*, Vol.11 (1978), 135-143.
19. M. Saigo, A certain boundary value problem for the Euler-Darboux equation I, *Math. Japonica*, Vol. 24 (4) (1979), 377-385.
20. A.A .Kilbas, H.M. Srivastava and J.J. Trujillo, Theory and applications of fractional differential equations, Elsevier, Amsterdam (2006).
21. S.G. Samko, A.A. Kilbas and O.I. Marichev, Fractional Integrals and Derivatives, Theory and Applications, Gordon and Breach, New York (1993).
22. E.M. Wright, The Asymptotic Expansion of the Generalized Hyper geometric function, *J. London Math. Soc.*, Vol.10 (1935), 286-293.
23. A.A. Kilbas, Fractional Calculus of the generalized Wright function, *Fract. Calc. Appl. Anal.*, Vol.8 (2), (2005), 113-126.
24. A.A. Kilbas and N. Sebastain, Generalized fractional integration of Bassel Function of first kind, *Integral Transforms and Spec. Funct.*, Vol.19 (12) (2008), 869- 883.
25. A.M. Mathai, A pathway to matrix-variate gamma and normal densities, *Linear Algebra and its applications*, 396 (2005), 317-328.
26. A.M. Mathai and H.J. Haubald, On generalized distribution and pathway, *Physics Letters*, 372 (2008), 2109-2113.

27. A.M. Mathai and H.J. Haubald, Pathway models, Super statistics, Trallis statistics and a generalized measure of entropy, *physica A* , 375 (2007), 110-122.
28. H.M. Srivastava and R.K. Saxena, operators of fractional integration and their application, *Appl. Math. Comput.* Vol. 118 (2001), 1-52.
29. P.Agarwal, G.V. Milovanovic and K.S. Nisar, A fractional integral operator involving the Mittag- Leftler type function with four parameters, *ser. Math. Inform.* , Vol.30 (5) (2015), 597-605.
30. K.S. Nisar, S.D. Purohit, M.S. Abouzaid, M.Al Qurashi and D. Baleanu, Generalized K-Mittag-Leffler function and it's composition withj pathway integral operators , *J. Nonlinear Sci. Appl.*, Vol.9 (2016), 3519-3526.
31. K.S. Nisar , A.F. Eata, M.Al-Dhaifallah and J. Choi , fractional calculus of generalized k-Mittag- Leftler function and it's application to statistical distribution , *Advances in Difference Equation Springer* (2016) , 1-17.
32. T.R. Prabhakar, A singular integral equation with a generalized Mittag-Leffler function in the kernel, *Yokohama Math. J.*, Vol. (19) (1977), 7-15.